

Real-Time PCR Assay for Rapid Detection of Epidemiologically and Clinically Significant *Mycobacterium tuberculosis* Beijing Genotype Isolates

Igor Mokrousov,^a Anna Vyazovaya,^a Viacheslav Zhuravlev,^b Tatiana Otten,^b Julie Millet,^c Wei-Wei Jiao,^d A-Dong Shen,^d Nalin Rastogi,^c Boris Vishnevsky,^b Olga Narvskaya^a

Laboratory of Molecular Microbiology, St. Petersburg Pasteur Institute, St. Petersburg, Russia^a; Laboratory of Etiological Diagnostics, Research Institute of Phthisiopulmonology, St. Petersburg, Russia^b; WHO Supranational TB Reference Laboratory, Unité de la Tuberculose et des Mycobactéries, Institut Pasteur de Guadeloupe, Abymes Cedex, Guadeloupe, France^c; National Key Discipline of Pediatrics, Key Laboratory of Major Diseases in Children (Capital Medical University), Ministry of Education, Beijing Pediatric Research Institute, Beijing Children's Hospital, Beijing, China^d

***Mycobacterium tuberculosis* Beijing genotype strains are rapidly disseminating, frequently hypervirulent, and multidrug resistant. Here, we describe a method for their rapid detection by real-time PCR that targets the specific IS6110 insertion in the *dnaA-dnaN* genome region. The method was evaluated with a geographically and genetically diverse collection representing areas in East Asia and the former Soviet Union in which the Beijing genotype is endemic and epidemic (i.e., major foci of its global propagation) and with clinical specimens.**

Mycobacterium tuberculosis is the major killer of humans in terms of morbidity and mortality. A closer look at its population structure reveals that different strains (lineages) of *M. tuberculosis* differ in their pathogenic capacities. The *M. tuberculosis* Beijing genotype is globally widespread and considered to be a fast-propagated family within *M. tuberculosis* (see references 1–3 and references therein). In Russia, the Beijing genotype strains are marked by increased virulence, as shown in a macrophage model (4), and are associated with multidrug resistance (5, 6). A “lethal combination” of the Beijing genotype and the human DC-SIGN–336G allele has been suggested for the male subgroup in the Siberian Slavic population (7). The capacity of the Beijing strains to escape a protection effect of the *Mycobacterium bovis* BCG vaccination in mice (8) suggests the clinical relevance of the early detection of this genotype.

For the above reasons, the availability of a simple method for detecting these strains is important for early adequate treatment and for epidemiological monitoring of their circulation. While no single standardized method for detecting the Beijing genotype has been approved, the availability of a wide array of methods would be helpful in view of the different technical capacities in different laboratories.

Previously, we reported the utility of the IS6110-based inverse PCR method for detecting Beijing strains based on specific and easily recognizable double-band profiles in agarose gel (9). One of these bands represents an IS6110 insertion in the *dnaA-dnaN* region in the proximity of *oriC* (position 1592 in the H37Rv genome [our unpublished data]), also previously described by Kurepina et al. (10). Here, we developed a method to detect this Beijing genotype-specific insertion, *dnaA-dnaN*::IS6110, in a real-time PCR format and evaluated it on a geographically and genetically diverse collection of 724 isolates from the main areas across Eurasia (Russia, Belarus, Kazakhstan, China, Vietnam, Japan) in which the Beijing genotype is epidemic and endemic. In addition, the method was preliminarily tested on DNA extracted from clinical specimens.

Real-time PCR with three primers and two labeled probes was performed in a single tube in multiplex format targeting the same

genomic region and detecting its two alleles, *dnaA-dnaN*::IS6110 specific for the Beijing genotype and the intact *dnaA-dnaN* region. Under this design, a PCR product should always be amplified with either one or another pair of primers and should consequently give rise to one or another fluorescent signal indicating either the Beijing or another genotype. The interpretation of the assay result is based on comparison of the two fluorescence curves for the same isolate, i.e., presence of a signal in one channel and the absence of a signal in another channel.

DNA was extracted from a bacterial culture using a recommended method (11). Purified DNA (0.5 to 1 ng) was added to the PCR mix (final volume, 25 μ l) containing 1.5 mM MgCl₂, 1 U *TaqF* hot-start DNA polymerase (InterLabService, Russia), 200 μ M (each) deoxynucleoside triphosphates (dNTP), 4 pmol each of the primers BGR (5'-CGCCGGGACTGTATGAGTCT), BGF2 (5'-CTCTCCCAGGTCACACCAGTCA), and BGRi (5'-TCGATGAACCACCTGACATGAC), and the probes BGPi (FAM-5'-CGGCATGTCCGGAGACTCCAGTTC [where FAM is 6-carboxy-fluorescein], 1.5 pmol) and BGP2 (HEX-5'-TGGCTGTGAGTGTGCTGTGCACA [where HEX is hexachloro fluorescein], 2.5 pmol). The PCR was run in a Rotor-Gene 6000 (Corbett Research) under the following conditions: 95°C for 10 min, 30 cycles at 94°C for 15 s, and 60°C for 50 s. Signal detection was performed at 60°C. A Beijing-specific insertion, *dnaA-dnaN*::IS6110, was revealed using the primers BGF2 and BGRi and the probe BGPi (green detection channel, wavelength of 510 nm). The intact

Received 14 November 2013 Returned for modification 30 December 2013

Accepted 3 February 2014

Published ahead of print 12 February 2014

Editor: S. A. Moser

Address correspondence to Igor Mokrousov, igormokrousov@yahoo.com.

Supplemental material for this article may be found at <http://dx.doi.org/10.1128/JCM.03193-13>.

Copyright © 2014, American Society for Microbiology. All Rights Reserved.

doi:10.1128/JCM.03193-13

dnaA-dnaN region was revealed using the primers BGF2 and BGR and the probe BGP2 (yellow detection channel, wavelength of 555 nm).

The Latin-American Mediterranean (LAM) family of *M. tuberculosis* was detected by a MboII PCR-restriction fragment length polymorphism (RFLP) analysis of *Rv0129c* 309 G→A mutation (12). The East-African Indian (EAI) family of *M. tuberculosis* was detected based on PCR detection of the RD239 deletion (13). To negate the possibility of a “pseudo-Beijing” genotype (14), the Beijing genotype isolates used in the initial optimization step in this study were confirmed to belong to the Beijing genotype by RD207 deletion analysis (15).

DNA from clinical specimens obtained from tuberculosis patients was extracted using commercial kits employing magnetic sorbent-based technology (AmpliSens [InterLabService] or M-Sorb-TUB [Syntol, Moscow, Russia]). Real-time PCR analysis of these samples was done as described above with a modification: 20 µl of DNA was added to a total volume of 50 µl PCR mix, and the signal accumulation was evaluated within 50 cycles. PCR inhibition was checked using a control assay implemented in the commercial kit to detect *M. tuberculosis* complex (AmpliTub-RV; Syntol), while an internal control (artificial DNA fragment cloned in the pUC19 plasmid) was amplified and detected in another fluorescence channel. In clinical samples, the concentrations of DNA were estimated using control samples and a calibration assay implemented in the same kit.

The assay was optimized using DNA of non-Beijing strains (including the reference strains H37Rv and BCG) and of Beijing strains of *M. tuberculosis* previously characterized using spoligotyping and confirmed to belong to the Beijing genotype by presence of the RD207 deletion.

Accumulation of the Beijing-specific signal was registered in the channel FAM/green (Fig. 1A), and accumulation of the non-Beijing genotype signal was registered in the channel HEX/yellow (Fig. 1B). The presence of Beijing DNA in a given sample was manifested as an exponential increase of the Beijing channel FAM signal and with the complete lack of signal in the non-Beijing channel HEX.

The optimized method was further evaluated with a representative and diverse (based on IS6110-RFLP and/or 24-loci mycobacterial interspersed repetitive-unit-variable-number tandem-repeat [MIRU-VNTR] typing and spoligotyping) collection of 724 DNA samples of *M. tuberculosis* isolates from the main areas of Beijing genotype endemicity in the former Soviet Union (Russia, Belarus, Kazakhstan) and East Asia (China, Vietnam, Japan) previously characterized by spoligotyping, IS6110-RFLP, and/or 24-MIRU-VNTR typing (see Table S1 and S2 in the supplemental material). This analysis revealed a complete concordance of the two methods in detecting the Beijing strain (real-time PCR and spoligotyping), with a few exceptions detailed below. Four hundred sixty-two isolates with a complete nine-signal spoligoprofile or abridged Beijing-like profiles with deleted signals were correctly identified as the Beijing genotype. In turn, 252 non-Beijing strains (genetic families LAM [*n* = 96], Haarlem [*n* = 26], Ural [*n* = 20], T [*n* = 66], X [*n* = 2], EAI [*n* = 25], unknown family [*n* = 17]) were correctly identified as non-Beijing.

The discrepant cases, showing the simultaneous presence of Beijing and non-Beijing real-time PCR signals, concerned 9 isolates from the former Soviet Union countries and 1 isolate from Japan with non-Beijing spoligotypes. A closer look at their spoli-




FIG 1 Fluorescence curves of a real-time PCR assay using DNA extracted from cultured bacteria. (A) Beijing genotype-specific signal (FAM channel, 510 nm); (B) other genotype-specific signal (HEX channel, 555 nm).

goprofiles revealed the presence of the 9 last signals characteristic of the Beijing genotype. Accordingly, these DNA samples were tested for additional markers that define other *M. tuberculosis* families. As a result, it was found that 9 isolates from countries in the former Soviet Union presented a mix of the LAM and Beijing genotypes, while 1 isolate from Japan presented a mix of the EAI and Beijing genotypes.

One hundred DNA samples extracted from clinical material (73 sputum samples, 8 bronchoalveolar lavage fluid [BALF] samples, and 19 other specimens, such as lung, tuberculoma, granulation, or chest cavity tissue) were randomly selected from the retrospective DNA collection. The concentration of DNA in the clinical specimens was estimated with a commercial kit to detect *M. tuberculosis* complex and was found to range from 4×10^2 to 1×10^6 copies per 10 µl. These specimens were subjected to the above-described real-time PCR under an increased number of cycles (*n* = 50). Clear-cut results were obtained for 86 of 100 samples, discriminating between Beijing (*n* = 59) and other genotypes (*n* = 27). PCR inhibition was checked using a control assay; as a result, no PCR inhibition was detected (data not shown). Thus, PCR failure (in 10 sputum and 4 BALF specimens) might have been due to the insufficient quantity/quality of the *M. tuberculosis* DNA.

A detailed discussion of different available methods for Beijing genotype detection (15–19) is beyond the scope of this short report. A larger prospective study should exploit the capacity of the developed method to identify the Beijing genotype in different kinds of clinical material. As a whole, a large-scale multicenter study is warranted to evaluate the different available PCR methods for detecting the Beijing genotype and for selecting those most suitable in terms of cost-effectiveness, performance, sensitivity, and specificity.

ACKNOWLEDGMENTS

We thank Ho Minh Ly, Nguen Ngoc Lan, Bahytul Zhakipbaeva, and Natalia Vasilenko for providing some of the mycobacterial DNA samples.

This work was partly supported by grants from the European Union's Seventh Framework Programme (FP7/2007–2013, grant agreement 261378), the Russian Foundation for Basic Research (grant 11-04-91172 GFEN_a), and the National Natural Science Foundation of China (grant 81071315) and by a research fellowship from the European Commission within the European Union's "Marie Curie Mobility" program to Igor Mokrousov (IIF contract 039389).

REFERENCES

- Bifani PJ, Mathema B, Kurepina NE, Kreiswirth BN. 2002. Global dissemination of the *Mycobacterium tuberculosis* W-Beijing family strains. *Trends Microbiol.* 10:45–52. [http://dx.doi.org/10.1016/S0966-842X\(01\)00277-6](http://dx.doi.org/10.1016/S0966-842X(01)00277-6).
- Demay C, Liens B, Burguière T, Hill V, Couvin D, Millet J, Mokrousov I, Sola C, Zozio T, Rastogi N. 2012. SITVITWEB—a publicly available international multimarker database for studying *Mycobacterium tuberculosis* genetic diversity and molecular epidemiology. *Infect. Genet. Evol.* 12:755–766. <http://dx.doi.org/10.1016/j.meegid.2012.02.004>.
- Mokrousov I. 2013. Insights into the origin, emergence, and current spread of a successful Russian clone of *Mycobacterium tuberculosis*. *Clin. Microbiol. Rev.* 26:342–360. <http://dx.doi.org/10.1128/CMR.00087-12>.
- Lasunskaja E, Ribeiro SC, Manicheva O, Gomes LL, Suffys PN, Mokrousov I, Ferrazoli L, Andrade MR, Kritski A, Otten T, Kipnis TL, da Silva WD, Vishnevsky B, Oliveira MM, Gomes HM, Baptista IF, Narvskaya O. 2010. Emerging multi-drug resistant *Mycobacterium tuberculosis* strains of the Beijing genotype circulating in Russia express a pattern of biological properties associated with enhanced virulence. *Microbes Infect.* 12:467–475. <http://dx.doi.org/10.1016/j.micinf.2010.02.008>.
- Drobniewski F, Balabanova Y, Nikolayevsky V, Ruddy M, Kuznetsov S, Zakharova S, Melentyev A, Fedorin I. 2005. Drug-resistant tuberculosis, clinical virulence, and the dominance of the Beijing strain family in Russia. *JAMA* 293:2726–2731. <http://dx.doi.org/10.1001/jama.293.22.2726>.
- Mokrousov I, Vyazovaya A, Otten T, Zhuravlev V, Pavlova E, Tarashkevich L, Krishevich V, Vishnevsky B, Narvskaya O. 2012. *Mycobacterium tuberculosis* population in northwestern Russia: an update from Russian-EU/Latvian border region. *PLoS One* 7:e41318. <http://dx.doi.org/10.1371/journal.pone.0041318>.
- Ogarkov O, Mokrousov I, Sinkov V, Zhdanova S, Antipina S, Savilov E. 2012. "Lethal" combination of *Mycobacterium tuberculosis* Beijing genotype and human CD2093–36G allele in Russian male population. *Infect. Genet. Evol.* 12:732–736. <http://dx.doi.org/10.1016/j.meegid.2011.10.005>.
- Lopez B, Aguilar D, Orozco H, Burger M, Espitia C, Ritacco V, Barrera L, Kremer K, Hernandez-Pando R, Huygen K, van Soolingen D. 2003. A marked difference in pathogenesis and immune response induced by different *Mycobacterium tuberculosis* genotypes. *Clin. Exp. Immunol.* 133:30–37. <http://dx.doi.org/10.1046/j.1365-2249.2003.02171.x>.
- Mokrousov I, Otten T, Vyazovaya A, Limeschenko E, Filipenko M, Sola C, Rastogi N, Steklova L, Vishnevsky B, Narvskaya O. 2003. PCR based methodology for detecting multi-drug resistant strains of *Mycobacterium tuberculosis* Beijing family circulating in Russia. *Eur. J. Clin. Microbiol. Infect. Dis.* 22:342–348. <http://dx.doi.org/10.1007/s10096-003-0944-0>.
- Kurepina NE, Sreevatsan S, Plikaytis BB, Bifani PJ, Connell ND, Donnelly RJ, van Soolingen D, Musser JM, Kreiswirth BN. 1998. Characterization of the phylogenetic distribution and chromosomal insertion sites of five IS6110 elements in *Mycobacterium tuberculosis*: non-random integration in the *dnaA-dnaN* region. *Tuber. Lung Dis.* 79:31–42. <http://dx.doi.org/10.1054/tuld.1998.0003>.
- van Embden JDA, Cave MD, Crawford JT, Dale JW, Eisenach KD, Gicquel B, Hermans P, Martin C, McAdam R, Shinnik TM, Small P. 1993. Strain identification of *Mycobacterium tuberculosis* by DNA fingerprinting: recommendations for a standardized methodology. *J. Clin. Microbiol.* 31:406–409.
- Gibson AL, Huard RC, Gey van Pittius NC, Lazzarini LC, Driscoll J, Kurepina N, Zozio T, Sola C, Spindola SM, Kritski AL, Fitzgerald D, Kremer K, Mardassi H, Chitale P, Brinkworth J, Garcia de Viedma D, Gicquel B, Pape JW, van Soolingen D, Kreiswirth BN, Warren RM, van Helden PD, Rastogi N, Suffys PN, Lapa e Silva J, Ho JL. 2008. Application of sensitive and specific molecular methods to uncover global dissemination of the major RDRio sublineage of the Latin American-Mediterranean *Mycobacterium tuberculosis* spoligotype family. *J. Clin. Microbiol.* 46:1259–1267. <http://dx.doi.org/10.1128/JCM.02231-07>.
- Gagneux S, DeRiemer K, Van T, Kato-Maeda M, de Jong BC, Narayanan S, Nicol M, Niemann S, Kremer K, Gutierrez MC, Hilty M, Hopewell PC, Small PM. 2006. Variable host-pathogen compatibility in *Mycobacterium tuberculosis*. *Proc. Natl. Acad. Sci. U. S. A.* 103:2869–2873. <http://dx.doi.org/10.1073/pnas.0511240103>.
- Fenner L, Malla B, Ninet B, Dubuis O, Stucki D, Borrell S, Huna T, Bodmer T, Egger M, Gagneux S. 2011. "Pseudo-Beijing": evidence for convergent evolution in the direct repeat region of *Mycobacterium tuberculosis*. *PLoS One* 6:e24737. <http://dx.doi.org/10.1371/journal.pone.0024737>.
- Tsolaki AG, Gagneux S, Pym AS, Goguet de la Salmoniere YO, Kreiswirth BN, van Soolingen D, Small PM. 2005. Genomic deletions classify the Beijing/W strains as a distinct genetic lineage of *Mycobacterium tuberculosis*. *J. Clin. Microbiol.* 43:3185–3191. <http://dx.doi.org/10.1128/JCM.43.7.3185-3191.2005>.
- Leung ET, Zheng L, Wong RY, Chan EW, Au TK, Chan RC, Lui G, Lee N, Ip M. 2011. Rapid and simultaneous detection of *Mycobacterium tuberculosis* complex and Beijing/W genotype in sputum by an optimized DNA extraction protocol and a novel multiplex real-time PCR. *J. Clin. Microbiol.* 49:2509–2515. <http://dx.doi.org/10.1128/JCM.00108-11>.
- Kurepina N, Likhoshvay E, Shashkina E, Mathema B, Kremer K, van Soolingen D, Bifani P, Kreiswirth BN. 2005. Targeted hybridization of IS6110 fingerprints identifies the W-Beijing *Mycobacterium tuberculosis* strains among clinical isolates. *J. Clin. Microbiol.* 43:2148–2154. <http://dx.doi.org/10.1128/JCM.43.5.2148-2154.2005>.
- Rindi L, Lari N, Garzelli C. 2010. A duplex real-time PCR assay for detection of *mutT4* and *mutT2* mutations in *Mycobacterium tuberculosis* of Beijing genotype. *J. Microbiol. Meth.* 81:203–204. <http://dx.doi.org/10.1016/j.mimet.2010.03.004>.
- Nakajima C, Tamaru A, Rahim Z, Poudel A, Maharjan B, Khin Saw Aye Ling H, Hattori T, Iwamoto T, Fukushima Y, Suzuki H, Suzuki Y, Matsuba T. 2013. Simple multiplex PCR assay for identification of Beijing family *Mycobacterium tuberculosis* isolates with a lineage-specific mutation in *Rv0679c*. *J. Clin. Microbiol.* 51:2025–2032. <http://dx.doi.org/10.1128/JCM.03404-12>.